

X-GAL

Shipping: On Dry/Blue Ice Catalog number
Exp. Date: See vial BIO-37035 : 1g
Batch No.: See vial



A Meridian Life Science® Company

Store at -20°C

Storage and stability: X-GAL should be stored at -20°C upon receipt. Store protected from light. When stored under optimum conditions, it is stable for up to 12 months from date of purchase.

Safety precautions: Harmful if swallowed. Irritating to eyes and skin. Please refer to the material safety data sheet for further information and safety handling practice.



Xn: Harmful

Notes:

For research use only
This product insert is a declaration of analysis at the time of manufacture

Description

5-bromo-4-chloro-3-indolyl β -D-galactopyranoside (X-GAL) is a chromogenic substrate for β -Galactosidase that forms an intense blue precipitate. It can be used in molecular biology to detect the *gal* gene product, and also in microbiology where it is used to detect micro-organisms which have β -Galactosidase activity (usually coliforms). It can be combined with the R-substrates to differentiate between two species of organisms on the same plate. X-GAL is soluble in N, N-dimethylformamide or dimethylsulfoxide (DMSO). Together with appropriate bacterial cloning vectors, host strains, and IPTG, X-GAL provides an easy way to distinguish between positive and negative clones after transformation.

Features

- Extremely pure, 99.5% by TLC (Thin Layer Chromatography)
- Intense blue precipitate upon hydrolysis

Applications

- Blue/White colony screening to distinguish recombinant (white) and non-recombinant (blue) colonies
- Immunoblotting
- Detect β -galactosidase activity in immunological and histochemical assays
- Microbiology and cell culture media

Protocol

Prepare a 20 mg/ml stock solution of X-Gal in N,N-dimethylformamide (DMF) or dimethylsulfoxide (DMSO). Sterilization is not required. Store stock solution in glass container protected from light at -20°C. Solutions may be stored at -20°C for 6-12 months. If a solution turns pink, it should be discarded. DMF dissolves some plastic materials. Direct addition of DMF containing solution to plastic Petri dishes should be avoided.

Procedures

Blue/White Colony Screening

a) X-Gal included in agar: Add 5 ml of X-Gal stock solution and 5 ml of 0.1 M isopropyl-b-D-thiogalactoside (IPTG, BIO-37082) for each 1L of autoclaved media agar (e.g. LB agar) containing appropriate antibiotics just before pouring. The media should be below 55°C. Plate cells on cooled agar and incubate overnight at 37°C.

b) X-Gal applied to top of agar: To a premade LB agar plate (e.g. prepared using LB agar), add 40 μ l of X-Gal stock solution (at room temperature) and 4 μ l of a 200 mg/ml solution of IPTG.

Spread solution over the entire surface of the plate. Incubate at 37°C until the fluid is no longer visible. This may take several hours. Plate cells and incubate overnight at 37°C.

Product Citations

1. Wilson, A.C., *et al. J. Bacteriol.* **190(15)**, 5522-5525(2008).
2. Liapis, E., *et al. NAR* **36(18)**, 5933-5945 (2008).
3. Corbett, D., *et al. J. Biol. Chem.* **282**, 33326-33335 (2007).
4. Toledo-Arana, A., *et al. J. Bacteriol.* **187(15)**, 5318-5329 (2005).
5. Staddon, J.H., *et al. Plasmid* **56(2)**, 102-111(2006).
6. Maruta, F., *et al. Can. Gene Therapy* **9(6)**, 543-552 (2002).
7. Marani, M., *et al. Mol. Cell. Biol.* **22(11)**, 3577-3589 (2002).

Associated Products

IPTG Solution	BIO-37082
IPTG Powder	BIO-37036
Quick-Stick Ligase	BIO-27027
T4 DNA Ligase	BIO-27026
SureClean	BIO-37042

Bioline Reagents Ltd
UNITED KINGDOM

Tel: +44(0)20 8830 5300
Fax: +44 (0)20 8452 2822

Bioline USA Inc.
USA

Tel: +1 508 880 8990
Fax: +1 508 880 8993

Bioline GmbH
GERMANY

Tel: +49(0)33 7168 1229
Fax: +49 (0)337168 1244

Bioline (Aust) Pty. Ltd
AUSTRALIA

Tel: +61 (0)2 9209 4180
Fax: +61 (0)2 9209 4763