

COA No: CA BN-0006

Version: 01

dATP 100mM

Storage Conditions:

-20°C

Lot number:

DA:013203

For Research Use Only

Expiry date: March 2015

Quality Control Parameters

Analysis	Specification	Result
Functional	A 3Kb Lambda DNA fragment is amplified with a dilution series of dATP, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37° C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5×10^{-3} U DNase.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection $9.7x10^{-3}$ ng/µl RNase.	Passed

Authorised by Jade James

Europe Headquarters UK



COA No: CA BN-0007

Version: 01

dCTP 100mM

Storage Conditions:

-20°C

Lot number:

DC:913103

For Research Use Only

Expiry date: March 2015

Quality Control Parameters

Analysis	Specification	Result
Functional	A 3Kb Lambda DNA fragment is amplified with a dilution series of dATP, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5×10^{-3} U DNase.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection 9.7x10 ⁻³ ng/µl RNase.	Passed

Authorised by Jade James

Europe Headquarters UK

info.uk@bioline.com Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 Europe Germany

info.de@bioline.com Tel: +49 (0)3371 681 229 Fax: +49 (0)3371 681 244 <u>America</u>

<u>Australia</u>

info.us@bioline.com Tel: +1 508 880 8990 Fax: +1 508 880 8993 info.aust@bioline.com

Tel: +61 (0)2 9209 4180 Fax: +61 (0)2 9209 4763



COA No: CA BN-0008

Version: 01

dGTP 100mM

Storage Conditions:

-20°C

Lot number:

DG:013203

For Research Use Only

Expiry date: March 2015

Quality Control Parameters

Analysis	Specification	Result
Functional	A 3Kb Lambda DNA fragment is amplified with a dilution series of dATP, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5×10^{-3} U DNase.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection 9.7x10 ⁻³ ng/µl RNase.	Passed

Authorised by Jade James

Europe Headquarters UK

info.uk@bioline.com Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 Europe Germany

info.de@bioline.com Tel: +49 (0)3371 681 229 Fax: +49 (0)3371 681 244 <u>America</u>

<u>a</u>

info.us@bioline.com Tel: +1 508 880 8990 Fax: +1 508 880 8993

<u>Australia</u>



COA No: CA BN-0009

Version: 01

dTTP 100mM

Storage Conditions:

-20°C

Lot number:

DT:913103

For Research Use Only

Expiry date: March 2015

Quality Control Parameters

Analysis	Specification	Result
Functional	A 3Kb Lambda DNA fragment is amplified with a dilution series of dATP, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37° C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5×10^{-3} U DNase.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection $9.7x10^{-3}$ ng/µl RNase.	Passed

Authorised by Jade James

Europe Headquarters UK

Fax: +61 (0)2 9209 4763