

# ACCUZYME™ DNA Polymerase

Shipping: On Dry/Blue Ice Catalog numbers

Batch No.: See vial BIO-21051 : 250 units (100 µL)

Concentration: 2.5 u/µL BIO-21052 : 500 units (200 µL)

Store at -20°C



A Meridian Life Science® Company

## Storage and stability:

The ACCUZYME is shipped on dry/blue ice. On arrival store at -20 °C for optimum stability. Repeated freeze/thaw cycles should be avoided.

## Expiry:

When stored under the recommended conditions and handled correctly, full activity of the kit is retained until the expiry date on the outer box label.

## Safety precautions:

Please refer to the material safety data sheet for further information.

## Unit definition:

One unit is defined as the amount of enzyme that incorporates 10nmoles of dNTPs into acid-insoluble form in 30 minutes at 72 °C.

## Notes:

Research use only.

## Features

- Very high yield
- High fidelity
- Amplifies fragments up to 5 kb

## Description

ACCUZYME™ DNA polymerase is a thermostable enzyme possessing 5'-3' DNA polymerase and 3'-5' proofreading exonuclease activities, offering high fidelity, even with demanding applications. ACCUZYME produces blunt-ended amplicons of up to 5 kb in length. ACCUZYME is supplied with 10x Reaction Buffer containing Mg<sup>2+</sup>, which provides optimal final reaction conditions for most experiments. In order to allow optimization of reaction conditions, additional MgCl<sub>2</sub> is provided.

## Applications

- Ideal for ultra-high fidelity for subsequent cloning
- Blunt-end cloning
- Site-directed mutagenesis

## Components

	250 Units	500 Units
ACCUZYME DNA Polymerase	100 µL	200 µL
10x AccuBuffer	1.2 mL	2 x 1.2 mL
50 mM MgCl <sub>2</sub> Solution	1.2 mL	1.2 mL

### PCR Reaction Conditions (for a 50 µL reaction)

The following protocol is for a standard 50 µL reaction and can be used as a starting point for reaction optimization. Please refer to the Important Considerations and PCR Optimization section.

10x AccuBuffer	5 µL
50 mM MgCl <sub>2</sub> Solution	Optional
100 mM dNTP Mix (see below)	0.5 µL
Template	as required
Primers 20 µM each	1 µL
ACCUZYME DNA Polymerase 2.5 U/µL	1 µL
Water (ddH <sub>2</sub> O)	up to 50 µL

### PCR cycling conditions:

Step	Temp.	Time	Cycles
Initial Denaturation	95 °- 98 °C	3 min	1
Denaturation	95 °- 98 °C	15 s	25-35
Annealing*	55 °-60 °C	15 s	
Extension	72 °C	1.5 - 2 min/kb	

\*Annealing temperature is primer dependent.

The conditions above are intended as a guide only; conditions will vary from reaction to reaction and may need optimization.

### Important considerations and PCR optimization

The optimal conditions will vary from reaction to reaction and are dependent on the template/primers used.

**10x AccuBuffer:** The 10x AccuBuffer comprises of 600 mM Tris-HCl, 60 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 100 mM KCl, 20 mM MgSO<sub>4</sub>, pH 8.3 at 25 °C.

The Mg<sup>2+</sup> concentration in the 1x AccuBuffer is 2 mM, this is the optimum concentration for ACCUZYME for most PCR reactions and should only be adjusted if necessary.

**ACCUZYME DNA Polymerase:** We recommend starting with 1 µL (2.5 Units) of ACCUZYME in a 50 µL reaction.

**Primers:** Forward and reverse primers are generally used at the final concentration of 0.2-0.6 µM each. As a starting point, we recommend using 0.4 µM final concentration (*i.e.* 20 pmol of each primer per 50 µL reaction volume). Too high a primer concentration can reduce the specificity of priming, resulting in non-specific products.

When designing primers we recommend using primer-design software such as Primer3 (<http://frodo.wi.mit.edu/primer3>) or visual OMP™ (<http://dnasoftware.com>). Primers should have a melting temperature (T<sub>m</sub>) of approximately 60 °C.

**Template:** The amount of template in the reaction depends mainly on the type of DNA used. For templates with low structural complexity, such as plasmid DNA, we recommend using 50 pg-10 ng DNA per 50 µL reaction volume. For eukaryotic genomic DNA, we recommend a starting amount of 200 ng DNA per 50 µL reaction, this can be varied between 5 ng - 500 ng. It is important to avoid using template re-suspended in EDTA-containing solutions (*e.g.* TE buffer) since EDTA chelates free Mg<sup>2+</sup>.

## Troubleshooting Guide

Problem	Possible Cause	Recommendation
<b>No PCR product</b>	Missing component	- Check reaction set-up and volumes used
	Defective component	- Check the aspect and the concentrations of all components as well as the storage conditions. If necessary test each component individually in controlled reactions
	Enzyme concentration too low	- Increase enzyme quantity in 0.5 U (0.2 µL) increments
	Cycling conditions not optimal	- Decrease the annealing temperature - Run a temperature gradient to determine the optimal annealing temperature - Increase the extension time, especially if amplifying a long target - Increase the number of cycles
	Difficult template e.g. GC or AT-rich, or high level of secondary structure	- Increase initial denaturation time to 5 minutes - Increase denaturation time
<b>Smearing or Non-Specific products</b>	Excessive cycling	- Decrease the number of cycles
	Extension time too long	- Decrease the extension time
	Annealing temperature too low	- Increase the annealing temperature
	Primer concentration too high	- Decrease primer concentration
	Contamination	- Replace each component in order to find the possible source of contamination - Set-up the PCR reaction and analyze the PCR product in separated areas

## Technical Support

If the troubleshooting guide does not solve the difficulty you are experiencing, please contact your local distributor or our Technical Support with details of reaction setup, cycling conditions and relevant data.

Email: [tech@bioline.com](mailto:tech@bioline.com)

## Associated Products

Product Name	Pack Size	Cat. No.
dNTP Set	4 x 25 µmol	BIO-39025
dNTP Mix	500 µL	BIO-39028
ACCUZYME™ Mix	2 x 1.25 mL	BIO-25027

## TRADEMARKS

1. ACCUZYME is a Trademark of Bioline Reagents Ltd

## Product Citations

1. Kitazono, A.A. *Gene* doi:10.1016/j.gene.2011.06.006 (2011).
2. Batchelor, D.J. *et al. Am. J. Physiol.* **300**, R67-R75 (2011).
3. Chiang, C. *et al. J. Bacteriol.* **193**, 52-62 (2011).
4. Chin, G.L., *et al. Appl. Envir. Microbiol.* **77**, 3451-3460 (2011).
5. Cheng, C., *et al. Mol. Cell. Biol.* **31**, 983-997 (2011).
6. Chakrabarti, M., *et al. Virol. J.* **7**, **181** (2010).
7. Silvestrini, F., *et al. Mol. Cell. Prot.*, **9**, 1437-48 (2010).
8. Williamson, D. S., *et al. Appl. Microbiol. Biotechnol.* **88**, 143-153 (2010).
9. Johnson M., *et al. NAR* **37(14)**, e98 (2009).
10. Pacheco, A., *et al. Microbiol.* **155**, 2021-2028 (2009).

Bioline Reagents Ltd  
UNITED KINGDOM  
Tel: +44 (0)20 8830 5300  
Fax: +44 (0)20 8452 2822

Bioline USA Inc.  
USA  
Tel: +1 508 880 8990  
Fax: +1 508 880 8993

Bioline GmbH  
GERMANY  
Tel: +49 (0)337 168 1229  
Fax: +49 (0)3371 68 1244

Bioline (Aust) Pty. Ltd  
AUSTRALIA  
Tel: +61 (0)2 9209 4180  
Fax: +61 (0)2 9209 4763

Bioline France  
FRANCE  
Tel: +33 (0)1 42 56 04 40  
Fax: +33 (0)9 70 06 62 10

Meridian Bioscience Asia Pte Ltd  
SINGAPORE  
Tel: +65 6774 7196  
Fax: +65 6774 6441