

COA No: CA BEM-0003

Version: 01

## MangoTaq DNA Polymerase

For Research Use Only

Storage Conditions:

-20°C

Lot number:

BTLM-415206

Expiry date:

June 2017

### **Quality Control Parameters**

Analysis	Specification	Result
Functional	Fragment of size 3Kb is amplified with a dilution series of Taq DNA Polymerase, using standard conditions and 30 cycles. Fragment of size 800bp is amplified with a dilution series of human genomic DNA, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5 x 10 <sup>-3</sup> U DNase.	Passed

Authorised by Jade James

Europe Headquarters UK

info.uk@bioline.com Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 Europe Germany

info.de@bioline.com Tel: +49 (0)3371 681 229 Fax: +49 (0)3371 681 244 <u>America</u>

America

info.us@bioline.com Tel: +1 508 880 8990 Fax: +1 508 880 8993 <u>Australia</u>

info.aust@bioline.com Tel: +61 (0)2 9209 4180 Fax: +61 (0)2 9209 4763



COA No: CA BB-0008

Version: 01

# **MangoTaq Buffer Colored**

For Research Use Only

**Storage Conditions:** 

-20°C

Lot number:

NHMR-515106

Expiry date:

June 2017

### **Quality Control Parameters**

Analysis	Specification	Result
Functional	Fragment of size 3000bp was amplified with a dilution series of Lamba DNA and a dilution series of MangoTaq™, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection $2.5 \times 10^{-3}$ U DNase.	Passed

Authorised by Jade James

CHOT CHIVE

Europe Headquarters UK



COA No: CA BB-0007

Version: 01

# **MangoTaq Buffer Colorless**

For Research Use Only

Storage Conditions:

-20°C

Lot number:

NHM-415106

Expiry date:

June 2017

### **Quality Control Parameters**

Analysis	Specification	Result
Functional	Fragment of size 3000bp was amplified with a dilution series of MangoTaq <sup>™</sup> , using standard conditions and 30 cycles. Fragment of size 800bp was amplified with a dilution series of Human Genomic DNA, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5 x 10 <sup>-3</sup> U DNase.	Passed

Authorised by Jade James

Europe Headquarters UK



COA No: CA BB-0014

Version: 01

# MgCl<sub>2</sub> Solution, 50mM

For Research Use Only

**Storage Conditions:** 

-20°C

Lot number:

MG-415106

Expiry date:

June 2017

### **Quality Control Parameters**

Analysis	Specification	Result
Functional	Fragments of sizes 800bp and 3000bp are amplified with a dilution series of BIOTAQ <sup>™</sup> DNA Polymerase, using standard conditions and 30 cycles. Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 2.5 x 10 <sup>-3</sup> U DNase.	Passed

Authorised by Jade James

CHOT CHIVE

